

A bi-endomorphism induces a new type of derivations on B-algebras

Thanatporn Bantaojai

Mathematics English Program

Faculty of Education

Valaya Alongkorn Rajabhat University under the Royal Patronage

Pathumtani 13180

Thailand

thanatporn.ban@vru.ac.th

Cholatis Suanoom

Science and Applied Science Center & Program of Mathematics

Faculty of Science and Technology

Kamphaeng Phet Rajabhat University

Kamphaeng Phet 62000

Thailand

cholatis.suanoom@gmail.com

Jirayu Phuto

Department of Mathematics

Faculty of Science

Naresuan University

Phitsanulok 65000

Thailand

jirayup60@email.nu.ac.th

Aiyared Iampan*

Fuzzy Algebras and Decision-Making Problems Research Unit

Department of Mathematics

School of Science

University of Phayao

Phayao 56000

Thailand

aiyared.ia@up.ac.th

Abstract. The goal of this paper is to introduce the concept of an (l, r) and an (r, l) - τ -derivation on a B-algebra which is induced by a left and a right bi-endomorphism and to provide important properties. The study found that the composition of (l, r) and an (r, l) - τ -derivations is also an (l, r) and an (r, l) - τ -derivation on a 0-commutative B-algebra, respectively. In addition, the relationship among those derivations is also considered.

Keywords: B-algebra, bi-endomorphism, derivation.

*. Corresponding author

1. Introduction

The notion of B-algebras was introduced by Neggers and Kim [10] in 2002, it is a new algebraic structure, they took some properties from BCI and BCH-algebras (see, [5, 7]), called a B-algebra.

In 2010, Al-Shehrie [3] introduced the notion of a left-right and a right-left derivation of a B-algebra and investigated some related properties. Also, he studied the notion of derivation of 0-commutative B-algebra and investigated some of its properties. Ardekani and Davvaz [4] introduced a generalization of a derivation of a B-algebra, that is, the notion of an f -derivation and an (f, g) -derivation of a B-algebra and investigated some properties of an (f, g) -derivation of a commutative B-algebra in 2014. In a UP-algebra, Sawika et al. [11] introduced the concepts of an (l, r) and an (r, l) -derivation and a derivation, Iampan [6] introduced the concept of an f -derivation, and in the next year, Tippanya et al. [12] introduced the concepts of a left and a right- f -derivation of type I, and of a left and a right- f -derivation of type II. In 2021, Muangkarn et al. [9] studied some properties of an outside and an inside f_q -derivation of a B-algebra. In addition, they defined and studied some properties of a (right-left) and a (left-right) f_q -derivation on a B-algebra from the concept of Al-Kadi [2].

In this paper, we introduce the concept of an (l, r) and an (r, l) - τ -derivation on a B-algebra which is induced by a left and a right bi-endomorphism and to provide important properties. In addition, the relationship among those derivations is also considered. Also, using the concept of a derivation in past investigate some of its properties.

2. Preliminaries

First, let's review some basic definitions and theorems that are required in our work.

Definition 2.1 ([10]). *By a B-algebra we mean an algebra $B = (B, \rightsquigarrow, 0)$ of type $(2, 0)$ satisfying the followings:*

$$(B1) \quad (\forall x \in B)(x \rightsquigarrow x = 0),$$

$$(B2) \quad (\forall x \in B)(x \rightsquigarrow 0 = x),$$

$$(B3) \quad (\forall x, y, z \in B)((x \rightsquigarrow y) \rightsquigarrow z = x \rightsquigarrow (z \rightsquigarrow (0 \rightsquigarrow y))).$$

A non-empty subset S a B-algebra B is called a subalgebra of B if $x \rightsquigarrow y \in S$, for all $x, y \in S$.

After that a B-algebra $(B, \rightsquigarrow, 0)$ is denoted by B .

Example 2.1 ([9]). Let $B = \{0, 1, 2, 3\}$ with the Cayley table as follows:

\rightsquigarrow	0	1	2	3
0	0	2	1	3
1	1	0	3	2
2	2	3	0	1
3	3	1	2	0

Then, B is a B-algebra. Let $S = \{0, 3\}$. Then, S is a subalgebra of B .

Theorem 2.1 ([10]). *If B is a B-algebra, then*

- (B4) $(\forall x, y \in B)((x \rightsquigarrow y) \rightsquigarrow (0 \rightsquigarrow y) = x)$,
- (B5) $(\forall x, y, z \in B)(x \rightsquigarrow (y \rightsquigarrow z) = (x \rightsquigarrow (0 \rightsquigarrow z)) \rightsquigarrow y)$,
- (B6) $(\forall x, y \in B)(x \rightsquigarrow y = 0 \Rightarrow x = y)$,
- (B7) $(\forall x \in B)(0 \rightsquigarrow (0 \rightsquigarrow x) = x)$,
- (B8) $(\forall x, y, z \in B)(x \rightsquigarrow z = y \rightsquigarrow z \Rightarrow x = y)$,
- (B9) $(\forall x, y, z \in B)(z \rightsquigarrow x = z \rightsquigarrow y \Rightarrow x = y)$.

Theorem 2.2 ([10]). *An algebra B is a B-algebra if and only if it satisfies the followings:*

- (B1) $(\forall x \in B)(x \rightsquigarrow x = 0)$,
- (B7) $(\forall x \in B)(0 \rightsquigarrow (0 \rightsquigarrow x) = x)$,
- (B10) $(\forall x, y, z \in B)((x \rightsquigarrow z) \rightsquigarrow (y \rightsquigarrow z) = x \rightsquigarrow y)$,
- (B11) $(\forall x, y \in B)(0 \rightsquigarrow (x \rightsquigarrow y) = y \rightsquigarrow x)$.

Definition 2.2 ([8]). *A B-algebra B is said to be 0-commutative if*

$$(\forall x, y \in B)(x \rightsquigarrow (0 \rightsquigarrow y) = y \rightsquigarrow (0 \rightsquigarrow x)).$$

Example 2.2. In Example 2.1, we have B is 0-commutative.

Theorem 2.3 ([8]). *If B is a 0-commutative B-algebra, then*

- (B12) $(\forall x, y \in B)((0 \rightsquigarrow x) \rightsquigarrow (0 \rightsquigarrow y) = y \rightsquigarrow x)$,
- (B13) $(\forall x, y, z \in B)((z \rightsquigarrow y) \rightsquigarrow (z \rightsquigarrow x) = x \rightsquigarrow y)$,
- (B14) $(\forall x, y, z \in B)((x \rightsquigarrow y) \rightsquigarrow z = (x \rightsquigarrow z) \rightsquigarrow y)$,
- (B15) $(\forall x, y \in B)((x \rightsquigarrow (x \rightsquigarrow y)) \rightsquigarrow y = 0)$,
- (B16) $(\forall x, y, z, t \in B)((x \rightsquigarrow z) \rightsquigarrow (y \rightsquigarrow t) = (t \rightsquigarrow z) \rightsquigarrow (y \rightsquigarrow x))$,

(B17) $(\forall x, y, z \in B)((x \rightsquigarrow y) \rightsquigarrow z = x \rightsquigarrow (y \rightsquigarrow z)),$

(B18) $(\forall x, y \in B)(x \rightsquigarrow (x \rightsquigarrow y) = y).$

For a B-algebra B , we denote $x \wedge y = y \rightsquigarrow (y \rightsquigarrow x)$, for all $x, y \in B$.

We can refer to (B7) and (B18) as follows:

(B7) $(\forall x, y \in B)(x \wedge 0 = x),$

(B18) $(\forall x, y \in B)(x \wedge y = x)$ if B is a 0-commutative B-algebra.

Definition 2.3 ([3]). *A self-map d on B is called*

(1) *a (left-right)-derivation ((l, r)-derivation) of B if*

$$(\forall x, y \in B)(d(x \rightsquigarrow y) = (d(x) \rightsquigarrow y) \wedge (x \rightsquigarrow d(y))),$$

(2) *a (right-left)-derivation ((r, l)-derivation) of B if*

$$(\forall x, y \in B)(d(x \rightsquigarrow y) = (x \rightsquigarrow d(y)) \wedge (d(x) \rightsquigarrow y)),$$

(3) *a derivation of B if it is both an (l, r) and an (r, l)-derivation of B .*

Definition 2.4 ([3]). *A self-map d on B is said to be regular if $d(0) = 0$. Otherwise, d is said to be irregular.*

Example 2.3. In Example 2.1, we define a self-map d on B by

$$d(x) = \begin{cases} 3, & \text{if } x = 0 \\ 2, & \text{if } x = 1 \\ 1, & \text{if } x = 2 \\ 0, & \text{if } x = 3. \end{cases}$$

Then, d is a derivation of B , and we see that d is irregular.

Definition 2.5 ([1]). *A non-empty subset I of B is called an ideal of B if it satisfies the followings:*

(I1) $0 \in I,$

(I2) $(\forall x, y \in B)(x \rightsquigarrow y \in I, y \in I \Rightarrow x \in I).$

We known that every subalgebra of B is an ideal.

Example 2.4. In Example 2.1, let $I = \{0, 2, 3\}$. Then, I is an ideal of B .

Proposition 2.1 ([1]). *Let I be an ideal of B . If $x \in B$ and $y \in I$ such that $x \rightsquigarrow y = 0$, then $x \in I$.*

Proposition 2.2 ([1]). *Let I be an ideal of B and $x, y \in I$. Then, $x \rightsquigarrow (0 \rightsquigarrow y) \in I$.*

Proposition 2.3 ([1]). *If I is an ideal of B such that $0 \rightsquigarrow x = x$, for all $x \in I$, then I is a subalgebra of B .*

3. Main results

In this section, we introduce bi-endomorphism on B-algebra and prove some results of some results of bi-endomorphisms of B-algebra B and its derivations as follows:

Definition 3.1. A mapping $\tau : B \times B \rightarrow B$ is called

(1) a left bi-endomorphism on B if

$$(\forall x, y, z \in B)(\tau(x \rightsquigarrow y, z) = \tau(x, z) \rightsquigarrow \tau(y, z)),$$

(2) a right bi-endomorphism on B if

$$(\forall x, y, z \in B)(\tau(x, y \rightsquigarrow z) = \tau(x, y) \rightsquigarrow \tau(x, z)),$$

(3) a bi-endomorphism on B if it is a left and a right bi-endomorphism on B .

Throughout this section, we assume that τ_l and τ_r are a left and a right bi-endomorphism of B , respectively.

Let $x \in B$. By (B2), we have

$$\tau_l(x, x) \rightsquigarrow 0 = \tau_l(x, x) = \tau_l(x \rightsquigarrow 0, x) = \tau_l(x, x) \rightsquigarrow \tau_l(0, x)$$

and

$$\tau_r(x, x) \rightsquigarrow 0 = \tau_r(x, x) = \tau_r(x, x \rightsquigarrow 0) = \tau_r(x, x) \rightsquigarrow \tau_r(x, 0).$$

By (B9), we have

$$(B19) \quad (\forall x \in B)(\tau_l(0, x) = 0),$$

$$(B20) \quad (\forall x \in B)(\tau_r(x, 0) = 0).$$

Example 3.1. In Example 2.1, we define a mapping $\tau_1 : B \times B \rightarrow B$ by

$$\tau_1(x, y) = \begin{cases} 3, & \text{if } (x, y) \in \{(1, 0), (2, 0)\} \\ 0, & \text{otherwise.} \end{cases}$$

Then, τ_1 is a left bi-endomorphism on B .

Definition 3.2. A mapping $\tau : B \times B \rightarrow B$ is said to be symmetric if $\tau(x, y) = \tau(y, x)$, for all $x, y \in B$.

Remark 3.1. For any B -algebra, define a mapping $0 : B \times B \rightarrow B$ by $0(x) = 0$, for all $x \in B$, which is a bi-endomorphism on B . Every symmetric left (right) bi-endomorphism on B is a bi-endomorphism.

Definition 3.3. An ideal I of B is called a τ -ideal of B if $\tau(x, y) \in I$, for all $x, y \in I$.

Example 3.2. From Example 2.4 and Example 3.1, we have $I = \{0, 2, 3\}$ is an ideal of B and $\tau_1(x, y) \in I$, for all $x, y \in I$. Thus I is a τ_1 -ideal on B .

Next, we generalize derivations on B-algebras with a left (right) bi-endomorphism $\tau : B \times B \rightarrow B$ from the concept of the Definition 2.3.

Definition 3.4. Let $\tau : B \times B \rightarrow B$ be a left (right) bi-endomorphism on B . A self-map d_τ on B is called

(1) an (l, r) - τ -derivation of B if

$$(\forall x, y \in B)(d_\tau(x \rightsquigarrow y) = (d_\tau(x) \rightsquigarrow \tau(x, y)) \wedge (\tau(x, y) \rightsquigarrow d_\tau(y))),$$

(2) an (r, l) - τ -derivation of B if

$$(\forall x, y \in B)(d_\tau(x \rightsquigarrow y) = (\tau(x, y) \rightsquigarrow d_\tau(y)) \wedge (d_\tau(x) \rightsquigarrow \tau(x, y))),$$

(3) a τ -derivation of B if it is both an (l, r) - and an (r, l) - τ -derivation of B .

Example 3.3. In Example 2.1, we define $\tau = 0$ and $d_\tau : B \rightarrow B$ by $d_\tau(x) = 3$. It is obvious that τ is a bi-endomorphism on B . Then, d_τ is an (l, r) - τ -derivation of B .

Definition 3.5. Let d be a self-map on B . An ideal I of B is said to be d -invariant if $d(I) \subseteq I$.

Example 3.4. In Example 2.1 and Example 2.4, we have $I = \{0, 2, 3\}$ is an ideal of B . From Example 3.3, we have d_τ is an (l, r) - τ -derivation of B . It is obvious that $d_\tau(x) \in I$, for all $x \in I$, that is, $d_\tau(I) \subseteq I$. Hence, I is d_τ -invariant.

Theorem 3.1. Let d_{τ_l} be an (l, r) - τ_l -derivation on B . Then, d_{τ_l} is regular if and only if $d_{\tau_l}(0 \rightsquigarrow x) = 0$, for all $x \in B$.

Proof. Suppose that d_{τ_l} is regular. Let $x \in B$. Then

$$\begin{aligned} d_{\tau_l}(0 \rightsquigarrow x) &= (d_{\tau_l}(0) \rightsquigarrow \tau_l(0, x)) \wedge (\tau_l(0, x) \rightsquigarrow d_{\tau_l}(x)) \\ \text{(B19)} \quad &= (0 \rightsquigarrow 0) \wedge (0 \rightsquigarrow d_{\tau_l}(x)) \\ \text{(B1)} \quad &= 0 \wedge (0 \rightsquigarrow d_{\tau_l}(x)) \\ &= (0 \rightsquigarrow d_{\tau_l}(x)) \rightsquigarrow ((0 \rightsquigarrow d_{\tau_l}(x)) \rightsquigarrow 0) \\ \text{(B2)} \quad &= (0 \rightsquigarrow d_{\tau_l}(x)) \rightsquigarrow (0 \rightsquigarrow d_{\tau_l}(x)) \\ \text{(B1)} \quad &= 0. \end{aligned}$$

Conversely, suppose that $d_{\tau_l}(0 \rightsquigarrow x) = 0$, for all $x \in B$. Then

$$\begin{aligned} 0 &= d_{\tau_l}(0 \rightsquigarrow 0) \\ &= (d_{\tau_l}(0) \rightsquigarrow \tau_l(0, 0)) \wedge (\tau_l(0, 0) \rightsquigarrow d_{\tau_l}(0)) \\ \text{(B19)} \quad &= (d_{\tau_l}(0) \rightsquigarrow 0) \wedge (0 \rightsquigarrow d_{\tau_l}(0)) \\ \text{(B2)} \quad &= d_{\tau_l}(0) \wedge (0 \rightsquigarrow d_{\tau_l}(0)) \\ &= (0 \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow ((0 \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow d_{\tau_l}(0)). \end{aligned}$$

By (B6) and (B2), we get $(0 \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow 0 = 0 \rightsquigarrow d_{\tau_l}(0) = (0 \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow d_{\tau_l}(0)$. Using (B9), we have $d_{\tau_l}(0) = 0$. Hence, d_{τ_l} is regular. \square

Theorem 3.2. *Let d_{τ_r} be an (l, r) - τ_r -derivation on B . Then, d_{τ_r} is regular and $\tau_r(0, x) = 0$, for all $x \in B$ if and only if $d_{\tau_r}(0 \rightsquigarrow x) = 0$, for all $x \in B$.*

Proof. Suppose that d_{τ_r} is regular and $\tau_r(0, x) = 0$, for all $x \in B$. Let $x \in B$. Then

$$\begin{aligned}
 d_{\tau_r}(0 \rightsquigarrow x) &= (d_{\tau_r}(0) \rightsquigarrow \tau_r(0, x)) \wedge (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \\
 &= (0 \rightsquigarrow 0) \wedge (0 \rightsquigarrow d_{\tau_r}(x)) \\
 \text{(B1)} \quad &= 0 \wedge (0 \rightsquigarrow d_{\tau_r}(x)) \\
 &= (0 \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow ((0 \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow 0) \\
 \text{(B2)} \quad &= (0 \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow (0 \rightsquigarrow d_{\tau_r}(x)) \\
 \text{(B1)} \quad &= 0.
 \end{aligned}$$

Conversely, suppose that $d_{\tau_r}(0 \rightsquigarrow x) = 0$, for all $x \in B$. By (B1), we have $d_{\tau_r}(0) = d_{\tau_r}(0 \rightsquigarrow 0) = 0$. Thus d_{τ_r} is regular. Let $x \in B$. Then

$$\begin{aligned}
 0 &= d_{\tau_r}(0 \rightsquigarrow x) \\
 &= (d_{\tau_r}(0) \rightsquigarrow \tau_r(0, x)) \wedge (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \\
 &= (0 \rightsquigarrow \tau_r(0, x)) \wedge (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \\
 &= (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow ((\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow (0 \rightsquigarrow \tau_r(0, x))).
 \end{aligned}$$

By (B6) and (B2), we get $(\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow 0 = \tau_r(0, x) \rightsquigarrow d_{\tau_r}(x) = (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \rightsquigarrow (0 \rightsquigarrow \tau_r(0, x))$. Using (B9), we have $0 \rightsquigarrow \tau_r(0, x) = 0$. Using (B6) again, we obtain $\tau_r(0, x) = 0$. \square

Next, we give a necessary and sufficient condition for (r, l) - τ_l -derivation and (r, l) - τ_r -derivation to be regular with some properties.

Theorem 3.3. *Let d_{τ_l} be an (r, l) - τ_l -derivation on B . Then, d_{τ_l} is regular and $\tau_l(x, 0) = 0$, for all $x \in B$ if and only if $d_{\tau_l}(x) = 0$, for all $x \in B$.*

Proof. Suppose that d_{τ_l} is regular and $\tau_l(x, 0) = 0$, for all $x \in B$. Let $x \in B$. Then

$$\begin{aligned}
 \text{(B2)} \quad d_{\tau_l}(x) &= d_{\tau_l}(x \rightsquigarrow 0) \\
 &= (\tau_l(x, 0) \rightsquigarrow d_{\tau_l}(0)) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \\
 &= (0 \rightsquigarrow 0) \wedge (d_{\tau_l}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad &= 0 \wedge d_{\tau_l}(x) \\
 &= d_{\tau_l}(x) \rightsquigarrow (d_{\tau_l}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad &= d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(x) \\
 \text{(B1)} \quad &= 0.
 \end{aligned}$$

Conversely, suppose that $d_{\tau_l}(x) = 0$, for all $x \in B$. Then, d_{τ_l} is regular. Let $x \in B$. Then

$$\begin{aligned}
 & 0 = d_{\tau_l}(x) \\
 \text{(B2)} \quad & = d_{\tau_l}(x \rightsquigarrow 0) \\
 & = (\tau_l(x, 0) \rightsquigarrow d_{\tau_l}(0)) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \\
 & = (\tau_l(x, 0) \rightsquigarrow 0) \wedge (0 \rightsquigarrow \tau_l(x, 0)) \\
 \text{(B2)} \quad & = \tau_l(x, 0) \wedge (0 \rightsquigarrow \tau_l(x, 0)) \\
 & = (0 \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow ((0 \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow \tau_l(x, 0)).
 \end{aligned}$$

By (B6) and (B2), we get $(0 \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow 0 = 0 \rightsquigarrow \tau_l(x, 0) = (0 \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow \tau_l(x, 0)$. Using (B9), we have $\tau_l(x, 0) = 0$. □

Theorem 3.4. *Let d_{τ_r} be an (r, l) - τ_r -derivation on B . Then, d_{τ_r} is regular if and only if $d_{\tau_r}(x) = 0$, for all $x \in B$.*

Proof. Suppose that d_{τ_r} is regular. Let $x \in B$. Then

$$\begin{aligned}
 \text{(B2)} \quad & d_{\tau_r}(x) = d_{\tau_r}(x \rightsquigarrow 0) \\
 & = (\tau_r(x, 0) \rightsquigarrow d_{\tau_r}(0)) \wedge (d_{\tau_r}(x) \rightsquigarrow \tau_r(x, 0)) \\
 \text{(B20)} \quad & = (0 \rightsquigarrow 0) \wedge (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad & = 0 \wedge d_{\tau_r}(x) \\
 & = d_{\tau_r}(x) \rightsquigarrow (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad & = d_{\tau_r}(x) \rightsquigarrow d_{\tau_r}(x) \\
 \text{(B1)} \quad & = 0.
 \end{aligned}$$

The converse is obvious that d_{τ_r} is regular. □

By (B19), we define an (l, r) - τ_l -derivation d_{τ_l} (or (r, l) - τ_l -derivation) on B by $d_{\tau_l}(x) = \tau_l(x, 0)$, for all $x \in B$, and obtain the following proposition.

Proposition 3.1. *Let d_{τ_l} be an (l, r) - τ_l -derivation on B . If $d_{\tau_l}(x) = \tau_l(x, 0)$, for all $x \in B$, then d_{τ_l} is the zero self-map on B and d_{τ_l} is regular.*

Proof. Suppose that $d_{\tau_l}(x) = \tau_l(x, 0)$, for all $x \in B$. By (B19), we have $d_{\tau_l}(0) = \tau_l(0, 0) = 0$, that is, d_{τ_l} is regular. Let $x \in B$. Then

$$\begin{aligned}
 \text{(B2)} \quad & d_{\tau_l}(x) = d_{\tau_l}(x \rightsquigarrow 0) \\
 & = (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \wedge (\tau_l(x, 0) \rightsquigarrow d_{\tau_l}(0)) \\
 & = (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(x)) \wedge (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \\
 \text{(B1)} \quad & = 0 \wedge (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \\
 & = (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow ((d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow 0) \\
 \text{(B2)} \quad & = (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \rightsquigarrow (d_{\tau_l}(x) \rightsquigarrow d_{\tau_l}(0)) \\
 \text{(B1)} \quad & = 0.
 \end{aligned}$$

□

Proposition 3.2. *Let d_{τ_l} be an (r, l) - τ_l -derivation on B . If $d_{\tau_l}(x) = \tau_l(x, 0)$, for all $x \in B$, then $d_{\tau_l}(0 \rightsquigarrow x) = 0 \rightsquigarrow d_{\tau_l}(x)$, for all $x \in B$ and d_{τ_l} is regular.*

Proof. Suppose that $d_{\tau_l}(x) = \tau_l(x, 0)$, for all $x \in B$. By (B19), we have $d_{\tau_l}(0) = \tau_l(0, 0) = 0$, that is, d_{τ_l} is regular. Let $x \in B$. Then

$$\begin{aligned}
 d_{\tau_l}(0 \rightsquigarrow x) &= (\tau_l(0, x) \rightsquigarrow d_{\tau_l}(x)) \wedge (d_{\tau_l}(0) \rightsquigarrow \tau_l(0, x)) \\
 \text{(B19)} \quad &= (0 \rightsquigarrow d_{\tau_l}(x)) \wedge (0 \rightsquigarrow 0) \\
 \text{(B1)} \quad &= (0 \rightsquigarrow d_{\tau_l}(x)) \wedge 0 \\
 \text{(B7)} \quad &= 0 \rightsquigarrow d_{\tau_l}(x).
 \end{aligned}$$

□

By (B20), we define an (l, r) - τ_l -derivation d_{τ_l} (or (r, l) - τ_l -derivation) on B by $d_{\tau_r}(x) = \tau_r(0, x)$, for all $x \in B$, and obtain the following proposition.

Proposition 3.3. *Let d_{τ_r} be an (l, r) - τ_r -derivation on B . If $d_{\tau_r}(x) = \tau_r(0, x)$, for all $x \in B$, then $d_{\tau_r}(0 \rightsquigarrow x) = 0 \rightsquigarrow d_{\tau_r}(x)$, for all $x \in B$ and d_{τ_r} is regular.*

Proof. Suppose that $d_{\tau_r}(x) = \tau_r(0, x)$, for all $x \in B$. By (B20), we have $d_{\tau_r}(0) = \tau_r(0, 0) = 0$, that is, d_{τ_r} is regular. Let $x \in B$. Then

$$\begin{aligned}
 d_{\tau_r}(0 \rightsquigarrow x) &= (d_{\tau_r}(0) \rightsquigarrow \tau_r(0, x)) \wedge (\tau_r(0, x) \rightsquigarrow d_{\tau_r}(x)) \\
 &= (d_{\tau_r}(0) \rightsquigarrow d_{\tau_r}(x)) \wedge (d_{\tau_r}(x) \rightsquigarrow d_{\tau_r}(x)) \\
 \text{(B1)} \quad &= (d_{\tau_r}(0) \rightsquigarrow d_{\tau_r}(x)) \wedge 0 \\
 \text{(B7)} \quad &= d_{\tau_r}(0) \rightsquigarrow d_{\tau_r}(x) \\
 &= 0 \rightsquigarrow d_{\tau_r}(x).
 \end{aligned}$$

□

Proposition 3.4. *Let d_{τ_r} be an (r, l) - τ_r -derivation on B . If $d_{\tau_r}(x) = \tau_r(0, x)$, for all $x \in B$, then d_{τ_r} is the zero self-map on B and d_{τ_r} is regular.*

Proof. Suppose that $d_{\tau_r}(x) = \tau_r(0, x)$, for all $x \in B$. By (B20), we have $d_{\tau_r}(0) = \tau_r(0, 0) = 0$, that is, d_{τ_r} is regular. Let $x \in B$. Then

$$\begin{aligned}
 d_{\tau_r}(x) &= d_{\tau_r}(x \rightsquigarrow 0) \\
 &= (\tau_r(x, 0) \rightsquigarrow d_{\tau_r}(0)) \wedge (d_{\tau_r}(x) \rightsquigarrow \tau_r(x, 0)) \\
 \text{(B20)} \quad &= (0 \rightsquigarrow 0) \wedge (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad &= 0 \wedge d_{\tau_r}(x) \\
 &= d_{\tau_r}(x) \rightsquigarrow (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad &= d_{\tau_r}(x) \rightsquigarrow d_{\tau_r}(x) \\
 \text{(B1)} \quad &= 0.
 \end{aligned}$$

□

Theorem 3.5. *Let d_{τ_l} be a regular (r, l) - τ_l -derivation on B . Every τ_l -ideal of B is d_{τ_l} -invariant.*

Proof. Let B be an element in a τ_l -ideal I of B . Then

$$\begin{aligned}
 \text{(B2)} \quad & d_{\tau_l}(x) \rightsquigarrow 0 = d_{\tau_l}(x) \\
 \text{(B2)} \quad & = d_{\tau_l}(x \rightsquigarrow 0) \\
 & = (\tau_l(x, 0) \rightsquigarrow d_{\tau_l}(0)) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \\
 & = (\tau_l(x, 0) \rightsquigarrow 0) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \\
 \text{(B2)} \quad & = \tau_l(x, 0) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \\
 & = (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow ((d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)) \rightsquigarrow \tau_l(x, 0)) \\
 \text{(B10)} \quad & = d_{\tau_l}(x) \rightsquigarrow (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)).
 \end{aligned}$$

By (B9), we get $0 = d_{\tau_l}(x) \rightsquigarrow \tau_l(x, 0)$. Since I is a τ_l -ideal of B and by (B6), we have $d_{\tau_l}(x) = \tau_l(x, 0) \in I$. Hence, I is d_{τ_l} -invariant. \square

Theorem 3.6. *Let d_{τ_r} be a regular (r, l) - τ_r -derivation on B . Every ideal of B is d_{τ_r} -invariant. In particular, every τ_r -ideal of B is d_{τ_r} -invariant.*

Proof. Let B be an element in an ideal I on B . Then

$$\begin{aligned}
 \text{(B2)} \quad & d_{\tau_r}(x) = d_{\tau_r}(x \rightsquigarrow 0) \\
 & = (\tau_r(x, 0) \rightsquigarrow d_{\tau_r}(0)) \wedge (d_{\tau_r}(x) \rightsquigarrow \tau_r(x, 0)) \\
 \text{(B20)} \quad & = (0 \rightsquigarrow 0) \wedge (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad & = 0 \wedge d_{\tau_r}(x) \\
 & = d_{\tau_r}(x) \rightsquigarrow (d_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B2)} \quad & = d_{\tau_r}(x) \rightsquigarrow d_{\tau_r}(x) \\
 \text{(B1)} \quad & = 0 \in I.
 \end{aligned}$$

Hence, I is d_{τ_r} -invariant. \square

Next, we focus on composition of τ_l and τ_r -derivations.

Theorem 3.7. *Let d_{τ_r} and D_{τ_r} be (l, r) - τ_r -derivations on a 0-commutative B -algebra B . If $\tau_r(x, y) = y$, for all $x, y \in B$, then $d_{\tau_r} \circ D_{\tau_r}$ is also an (l, r) - τ_r -derivation on B .*

Proof. Suppose that $\tau_r(x, y) = y$, for all $x, y \in B$. Then

$$\begin{aligned}
 & (d_{\tau_r} \circ D_{\tau_r})(x \rightsquigarrow y) \\
 & = d_{\tau_r}(D_{\tau_r}(x \rightsquigarrow y)) \\
 & = d_{\tau_r}((D_{\tau_r}(x) \rightsquigarrow \tau_r(x, y)) \wedge (\tau_r(x, y) \rightsquigarrow D_{\tau_r}(y))) \\
 & = d_{\tau_r}((D_{\tau_r}(x) \rightsquigarrow y) \wedge (y \rightsquigarrow D_{\tau_r}(y)))
 \end{aligned}$$

$$\begin{aligned}
\text{(B18)} \quad &= d_{\tau_r}(D_{\tau_r}(x) \rightsquigarrow y) \\
&= (d_{\tau_r}(D_{\tau_r}(x)) \rightsquigarrow \tau_r(D_{\tau_r}(x), y)) \wedge (\tau_r(D_{\tau_r}(x), y) \rightsquigarrow d_{\tau_r}(y)) \\
\text{(B18)} \quad &= d_{\tau_r}(D_{\tau_r}(x)) \rightsquigarrow \tau_r(D_{\tau_r}(x), y) \\
&= d_{\tau_r}(D_{\tau_r}(x)) \rightsquigarrow y \\
&= d_{\tau_r}(D_{\tau_r}(x)) \rightsquigarrow \tau_r(x, y) \\
\text{(B18)} \quad &= (d_{\tau_r}(D_{\tau_r}(x)) \rightsquigarrow \tau_r(x, y)) \wedge (\tau_r(x, y) \rightsquigarrow d_{\tau_r}(D_{\tau_r}(y))) \\
&= ((d_{\tau_r} \circ D_{\tau_r})(x) \rightsquigarrow \tau_r(x, y)) \wedge (\tau_r(x, y) \rightsquigarrow (d_{\tau_r} \circ D_{\tau_r})(y)).
\end{aligned}$$

Hence, $d_{\tau_r} \circ D_{\tau_r}$ is an (l, r) - τ_l -derivation on B . \square

Theorem 3.8. *Let d_{τ_l} and D_{τ_l} be (r, l) - τ_l -derivations on a 0-commutative B -algebra B . If $\tau_l(x, y) = x$, for all $x, y \in B$, then $d_{\tau_l} \circ D_{\tau_l}$ is also an (r, l) - τ_l -derivation on B .*

Proof. Suppose that $\tau_r(x, y) = x$, for all $x, y \in B$. Then

$$\begin{aligned}
&(d_{\tau_l} \circ D_{\tau_l})(x \rightsquigarrow y) \\
&= d_{\tau_l}(D_{\tau_l}(x \rightsquigarrow y)) \\
&= d_{\tau_l}((\tau_l(x, y) \rightsquigarrow D_{\tau_l}(y)) \wedge (D_{\tau_l}(x) \rightsquigarrow \tau_l(x, y))) \\
&= d_{\tau_l}((x \rightsquigarrow D_{\tau_l}(y)) \wedge (D_{\tau_l}(x) \rightsquigarrow x)) \\
\text{(B18)} \quad &= d_{\tau_l}(x \rightsquigarrow D_{\tau_l}(y)) \\
&= (\tau_l(x, D_{\tau_l}(y)) \rightsquigarrow d_{\tau_l}(D_{\tau_l}(y))) \wedge (d_{\tau_l}(x) \rightsquigarrow \tau_l(x, D_{\tau_l}(y))) \\
\text{(B18)} \quad &= \tau_l(x, D_{\tau_l}(y)) \rightsquigarrow d_{\tau_l}(D_{\tau_l}(y)) \\
&= x \rightsquigarrow d_{\tau_l}(D_{\tau_l}(y)) \\
&= \tau_l(x, y) \rightsquigarrow d_{\tau_l}(D_{\tau_l}(y)) \\
\text{(B18)} \quad &= (\tau_l(x, y) \rightsquigarrow d_{\tau_l}(D_{\tau_l}(y))) \wedge (d_{\tau_l}(D_{\tau_l}(x)) \rightsquigarrow \tau_l(x, y)) \\
&= (\tau_l(x, y) \rightsquigarrow (d_{\tau_l} \circ D_{\tau_l})(y)) \wedge ((d_{\tau_l} \circ D_{\tau_l})(x) \rightsquigarrow \tau_l(x, y)).
\end{aligned}$$

Hence, $d_{\tau_l} \circ D_{\tau_l}$ is an (r, l) - τ_l -derivation on B . \square

Proposition 3.5. *Let d_{τ_r} be an (r, l) - τ_r -derivation and D_{τ_r} be an (l, r) - τ_r -derivation on a 0-commutative B -algebra B . Then*

- (1) $(\forall x \in B)((d_{\tau_r} \circ D_{\tau_r})(x) = 0)$ if d_{τ_r} is regular,
- (2) $(\forall x \in B)((D_{\tau_r} \circ d_{\tau_r})(x) = D_{\tau_r}(0) \rightsquigarrow \tau_r(0, d_{\tau_r}(0)))$,
- (3) $(\forall x \in B)((d_{\tau_r} \circ D_{\tau_r})(x) = 0 = (D_{\tau_r} \circ d_{\tau_r})(x))$ if d_{τ_r} and D_{τ_r} are regular.

Proof. (1) Suppose that d_{τ_r} is regular. Let $x \in B$. Then

$$\begin{aligned}
 \text{(B2)} \quad & (d_{\tau_r} \circ D_{\tau_r})(x) = (d_{\tau_r} \circ D_{\tau_r})(x \rightsquigarrow 0) \\
 & = d_{\tau_r}(D_{\tau_r}(x \rightsquigarrow 0)) \\
 \text{(B18)} \quad & = d_{\tau_r}(D_{\tau_r}(x) \rightsquigarrow \tau_r(x, 0)) \\
 \text{(B20)} \quad & = d_{\tau_r}(D_{\tau_r}(x) \rightsquigarrow 0) \\
 \text{(B18)} \quad & = \tau_r(D_{\tau_r}(x), 0) \rightsquigarrow d_{\tau_r}(0) \\
 \text{(B20)} \quad & = 0 \rightsquigarrow 0 \\
 \text{(B1)} \quad & = 0.
 \end{aligned}$$

(2) Let $x \in B$. Then

$$\begin{aligned}
 \text{(B2)} \quad & (D_{\tau_r} \circ d_{\tau_r})(x) = (D_{\tau_r} \circ d_{\tau_r})(x \rightsquigarrow 0) \\
 & = D_{\tau_r}(d_{\tau_r}(x \rightsquigarrow 0)) \\
 \text{(B18)} \quad & = D_{\tau_r}(\tau_r(x, 0) \rightsquigarrow d_{\tau_r}(0)) \\
 \text{(B20)} \quad & = D_{\tau_r}(0 \rightsquigarrow d_{\tau_r}(0)) \\
 \text{(B18)} \quad & = D_{\tau_r}(0) \rightsquigarrow \tau_r(0, d_{\tau_r}(0)).
 \end{aligned}$$

(3) It is straightforward by (1) and (2). □

Let $\text{Der}_{\tau}(B)$ be the set of all τ -derivations on a 0-commutative B-algebra B . For $d_{\tau}, D_{\tau} \in \text{Der}_{\tau}(B)$, we define the binary operation \wedge on $\text{Der}_{\tau}(B)$ as follows:

$$(\forall x \in B)((d_{\tau} \wedge D_{\tau})(x) = d_{\tau}(x) \wedge D_{\tau}(x)).$$

Indeed, let $x \in B$. Then

$$\begin{aligned}
 \text{(B18)} \quad & (d_{\tau} \wedge D_{\tau})(x) = d_{\tau}(x) \wedge D_{\tau}(x) \\
 & = d_{\tau}(x).
 \end{aligned}$$

Hence, $d_{\tau} \wedge D_{\tau} = d_{\tau}$, for all $d_{\tau}, D_{\tau} \in \text{Der}_{\tau}(B)$.

Therefore, we immediately get the result as the following proposition.

Proposition 3.6. *For a 0-commutative B-algebra B , $(\text{Der}_{\tau}(B), \wedge)$ is the left zero semigroup.*

4. Conclusion and discussion

In this paper, we have introduced the concept of an (l, r) and an (r, l) - τ -derivation on a B-algebra which is induced by a left and a right bi-endomorphism and provided important properties. The study found that the composition of (l, r) and an (r, l) - τ -derivations is also an (l, r) and an (r, l) - τ -derivation on a 0-commutative B-algebra, respectively. In addition, we can show that the set of all τ -derivations on a 0-commutative B-algebra B is the left zero semigroup. Finally, the study of a bi-endomorphism on other algebras (d/BH/BF/BG-algebras) is an interesting open problem.

Acknowledgment

The authors would like to thanks the Research and Development Institute, Kamphaeng Phet Rajabhat University to support.

References

- [1] H. K. Abdullah, A. A. Atshan, *Complete ideal and n -ideal of B -algebra*, Appl. Math. Sci., 11 (2017), 1705-1713.
- [2] D. Al-Kadi, *f_q -Derivations of G -algebra*, Int. J. Math. Comput. Sci., 2016 (2016), Article ID 9276096.
- [3] N. O. Al-Shehrie, *Derivation of B -algebras*, J. King Abdulaziz Univ. Sci., 22 (2010), 71-83.
- [4] L. K. Ardekani, B. Davvaz, *On (f, g) -derivations of B -algebras*, Mat. Vesnik, 66 (2014), 125-132.
- [5] Q. P. Hu, X. Li, *On BCH -algebras*, Math. Seminar Notes, 11 (1983), 313-320.
- [6] A. Iampan, *Derivations of UP -algebras by means of UP -endomorphisms*, Alg. Struc. Appl., 3 (2016), 1-20.
- [7] Y. B. Jun, X. L. Xin, *On derivations of BCI -algebras*, Inform. Sci., 159 (2004), 167-176.
- [8] H. S. Kim, H. G. Park, *On 0 -commutative B -algebras*, Sci. Math. Japon. Online, e-2005 (2005), 31-36.
- [9] P. Muangkarn, C. Suanoom, P. Pengyim, A. Iampan, *f_q -derivations of B -algebras*, J. Math. Comput. Sci., 11 (2021), 2047-2057.
- [10] J. Neggers, H. S. Kim, *On B -algebras*, Mat. Vesnik, 54 (2002), 21-29.
- [11] K. Sawika, R. Intasan, A. Kaewwasri, A. Iampan, *Derivations of UP -algebras*, Korean J. Math., 24 (2016), 345-367.
- [12] T. Tippanya, N. Iam-art, P. Moonfong, A. Iampan, *A new derivations of UP -algebras by means of UP -endomorphisms*, Algebra Lett., 2017 (2017), Article ID 4.

Accepted: July 14, 2021