

Tri-Endomorphisms on BCH-Algebras

Patchara Muangkarn¹, Cholatis Suanoom¹, Jirayu Phuto², Aiyared Iampan^{3,*}

¹*Science and Applied Science Center, Program of Mathematics, Kamphaeng Phet Rajabhat University, Kamphaeng Phet 62000, Thailand*

²*Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand*

³*Fuzzy Algebras and Decision-Making Problems Research Unit, Department of Mathematics, School of Science, University of Phayao, Mae Ka, Mueang, Phayao 56000, Thailand*

*Corresponding author: aiyared.ia@up.ac.th

Abstract. In this paper, we use the concept of endomorphisms and bi-endomorphisms as a model to create tri-endomorphisms on of BCH-algebras. We introduce the concepts of left tri-endomorphisms, central tri-endomorphisms, right tri-endomorphisms, and complete tri-endomorphisms of BCH-algebras and provide some properties. In addition, we obtain the properties between those tri-endomorphisms and some subsets of BCH-algebras.

1. Introduction

The algebraic structures of BCK-algebras and BCI-algebras were studied by Iséki and his colleague [4, 5]. In 1983, Hu and Li [3] generalized a new class of algebras from BCI-algebras, namely, a BCH-algebra. Next, Bandru and Rafi [1] introduced a new algebra, called a G -algebra. BCH-algebras are also being studied extensively later, [2, 3].

In this paper, we use the concept of endomorphisms and bi-endomorphisms as a model to create tri-endomorphisms. We introduce the concepts of left tri-endomorphisms, central tri-endomorphisms, right tri-endomorphisms, and complete tri-endomorphisms of BCH-algebras and provide some properties.

Before studying, we will review the definitions and well-known results.

Received: Jul. 31, 2022.

2020 *Mathematics Subject Classification.* 06F35, 03G25.

Key words and phrases. BCH-algebra; left tri-endomorphism; central tri-endomorphism; right tri-endomorphism; complete tri-endomorphism.

Definition 1.1. [3] A BCH-algebra is a non-empty set X with an element 0 and a binary operation $*$ satisfying the following conditions:

$$(BCH1) (\forall x \in X)(x * x = 0),$$

$$(BCH2) (\forall x, y \in X)(x * y = 0, y * x = 0 \Rightarrow x = y),$$

$$(BCH3) (\forall x, y, z \in X)((x * y) * z = (x * z) * y).$$

In a BCH-algebra $X = (X, *, 0)$, the binary relation \leq on X is defined as follows:

$$(\forall x, y \in X)(x \leq y \Leftrightarrow x * y = 0).$$

Example 1.1. Let $X = \{0, a, b, c\}$ with the following Cayley table as follows:

$*$	0	a	b	c
0	0	0	0	0
a	a	0	c	c
b	b	0	0	b
c	c	0	0	0

Then $X = (X, *, 0)$ is a BCH-algebra.

Proposition 1.1. [2,3] Let $X = (X, *, 0)$ be a BCH-algebra. Then the following hold: for all $x, y \in X$,

$$(BCH4) (\forall x, y \in X)(x * (x * y) \leq y),$$

$$(BCH5) (\forall x \in X)(x * 0 = 0 \Rightarrow x = 0),$$

$$(BCH6) (\forall x, y \in X)(0 * (x * y) = (0 * x) * (0 * y)),$$

$$(BCH7) (\forall x \in X)(x * 0 = x),$$

$$(BCH8) (\forall x, y \in X)((x * y) * x = 0 * y),$$

$$(BCH9) (\forall x, y \in X)(x \leq y \Rightarrow 0 * x = 0 * y).$$

For a BCH-algebra $X = (X, *, 0)$, some interesting subsets of X play a significant role in the investigation of its properties described below.

Definition 1.2. A non-empty subset Y of a BCH-algebra $X = (X, *, 0)$ is called a subalgebra of X if $x * y \in Y$ for all $x, y \in Y$. A non-empty subset I of a BCH-algebra $X = (X, *, 0)$ is called an ideal of X if

$$(1) 0 \in I,$$

$$(2) (\forall x, y \in X)(x * y \in I, x \in I \Rightarrow y \in I).$$

2. Main results

In this section, we introduce the concepts of left tri-endomorphisms, central tri-endomorphisms, right tri-endomorphisms, and complete tri-endomorphisms of BCH-algebras as follows.

Definition 2.1. Let $X = (X, *, 0)$ be a BCH-algebra. A mapping $f : X^3 \rightarrow X$ is called

- (1) a left tri-endomorphism on X if $(\forall w, x, y, z \in X)(f(x * w, y, z) = f(x, y, z) * f(w, y, z))$,
- (2) a central tri-endomorphism on X if $(\forall w, x, y, z \in X)(f(x, y * w, z) = f(x, y, z) * f(x, w, z))$,
- (3) a right tri-endomorphism on X if $(\forall w, x, y, z \in X)(f(x, y, z * w) = f(x, y, z) * f(x, y, w))$,
- (4) a complete tri-endomorphism on X if $(\forall a, b, c, x, y, z \in X)(f(x * a, y * b, z * c) = f(x, y, z) * f(a, b, c))$.

Example 2.1. In Example 1.1, we define $f_l : X^3 \rightarrow X$ by

$$f_l(x, y, z) = \begin{cases} x & \text{if } y = z = 0, \\ 0 & \text{otherwise.} \end{cases}$$

Then f_l is a left tri-endomorphism on X .

Proposition 2.1. Let $X = (X, *, 0)$ be a BCH-algebra and f_l be a left tri-endomorphism on X . Then

- (1) $(\forall y, z \in X)(f_l(0, y, z) = 0)$,
- (2) $(\forall w, x, y, z \in X)(x \leq w \Rightarrow f_l(x, y, z) \leq f_l(w, y, z))$.

Proof. (1) Let $y, z \in X$. Then, by BCH1, we have $f_l(0, y, z) = f_l(0 * 0, y, z) = f_l(0, y, z) * f_l(0, y, z) = 0$.

(2) Let $w, x, y, z \in X$ be such that $x \leq w$. Then, by (1), we have $0 = f_l(0, y, z) = f_l(x * w, y, z) = f_l(x, y, z) * f_l(w, y, z)$. Hence, $f_l(x, y, z) \leq f_l(w, y, z)$. \square

Similarly, the properties of central and right tri-endomorphisms are easily obtained.

Proposition 2.2. Let $X = (X, *, 0)$ be a BCH-algebra and f_c be a central tri-endomorphism on X .

Then

- (1) $(\forall x, z \in X)(f_c(x, 0, z) = 0)$,
- (2) $(\forall w, x, y, z \in X)(y \leq w \Rightarrow f_c(x, y, z) \leq f_c(x, w, z))$.

Proposition 2.3. Let $X = (X, *, 0)$ be a BCH-algebra and f_r be a right tri-endomorphism on X . Then

- (1) $(\forall x, y \in X)(f_r(x, y, 0) = 0)$,
- (2) $(\forall w, x, y, z \in X)(z \leq w \Rightarrow f_r(x, y, z) \leq f_r(x, y, w))$.

Theorem 2.1. Let $X = (X, *, 0)$ be a BCH-algebra and f be a complete tri-endomorphism on X .

Then

- (1) $f(0, 0, 0) = 0$,
- (2) if S is a subalgebra of X , then $f(S^3)$ is also a subalgebra of X ,
- (3) if S is an ideal of X and f is bijective, then $f(S^3)$ is also an ideal of X ,
- (4) if f is a left tri-endomorphism on X , then $f(x, y, z) * f(x, 0, 0) = 0$ for any $x, y, z \in X$,
- (5) if f is a central tri-endomorphism on X , then $f(x, y, z) * f(0, y, 0) = 0$ for any $x, y, z \in X$,
- (6) if f is a right tri-endomorphism on X , then $f(x, y, z) * f(0, 0, z) = 0$ for any $x, y, z \in X$,

(7) if f is a left and right (central and right, left and central) tri-endomorphism on X , then $f(x, y, z) = 0$ for any $x, y, z \in X$, i.e., f is the zero map.

Proof. (1) By BCH1, we have $f(0, 0, 0) = f(0 * 0, 0 * 0, 0 * 0) = f(0, 0, 0) * f(0, 0, 0) = 0$.

(2) Suppose that S is a subalgebra of X . Let $a, b \in f(S^3)$. Then there exist $(x_1, y_1, z_1), (x_2, y_2, z_2) \in S^3$ such that $a = f(x_1, y_1, z_1)$ and $b = f(x_2, y_2, z_2)$. Thus $a * b = f(x_1, y_1, z_1) * f(x_2, y_2, z_2) = f(x_1 * x_2, y_1 * y_2, z_1 * z_2) \in f(S^3)$. Hence, $f(S^3)$ is a subalgebra of X .

(3) Suppose that S is an ideal of X and f is bijective. Since $0 \in S$ and by (1), we have $0 = f(0, 0, 0) \in f(S^3)$. Assume that $x * y \in f(S^3)$ and $x \in f(S^3)$. There exist $(x_1, y_1, z_1), (x_2, y_2, z_2) \in S^3$ such that $x * y = f(x_1, y_1, z_1)$ and $x = f(x_2, y_2, z_2)$. Since f is surjective, there exists $(a, b, c) \in X^3$ such that $y = f(a, b, c)$. Thus $f(S^3) \ni f(x_1, y_1, z_1) = x * y = f(x_2, y_2, z_2) * f(a, b, c) = f(x_2 * a, y_2 * b, z_2 * c)$. Since f is injective, we have $x_2 * a, y_2 * b, z_2 * c \in S$. Since S is an ideal of X , we get $a, b, c \in S$. Thus $y = f(a, b, c) \in f(S^3)$. Hence, $f(S^3)$ is an ideal of X .

(4)-(6) It is obvious from Propositions 2.1-2.3.

(7) Suppose that f is a left and right tri-endomorphism on X . Let $x, y, z \in X$. Then, by Propositions 2.1 and 2.3, BCH1, BCH7 $0 = f(0, y, z) = f(x * x, y * 0, z * 0) = f(x, y, z) * f(x, 0, 0) = f(x, y, z) * 0 = f(x, y, z)$. Hence, f is the zero map on X . \square

Let $T_l(X)$ (resp., $T_c(X)$, $T_r(X)$ and $T(X)$) be the set of all left tri-endomorphisms (resp., right, central and complete tri-endomorphisms) on a BCH-algebra $X = (X, *, 0)$. We define an operation \star on $T_l(X)$ by $(\forall x, y, z \in X)((f \star g)(x, y, z) = f(x, y, z) * g(x, y, z))$. Let $f \in T_l(X)$ and $x, y, z \in X$. Then $(f \star f)(x, y, z) = f(x, y, z) * f(x, y, z) = 0$. This means that $f \star f = 0_X$, where $0_X : X^3 \rightarrow X$ is a function that maps all members to 0. Let $f, g \in T_l(X)$ be such that $f \star g = 0_X$ and $g \star f = 0_X$. Then for all $x, y, z \in X$, $0 = (f \star g)(x, y, z) = f(x, y, z) * g(x, y, z)$ and $0 = (g \star f)(x, y, z) = g(x, y, z) * f(x, y, z)$. Since $g(x, y, z), f(x, y, z) \in X$, we have $f(x, y, z) = g(x, y, z)$ for all $x, y, z \in X$. Hence, $f = g$. Let $f, g, h \in T_l(X)$ and $x, y, z \in X$. Then $((f \star g) \star h)(x, y, z) = (f \star g)(x, y, z) * h(x, y, z) = (f(x, y, z) * g(x, y, z)) * h(x, y, z) = (f(x, y, z) * h(x, y, z)) * g(x, y, z) = (f \star h)(x, y, z) * g(x, y, z) = ((f \star h) \star g)(x, y, z)$. Hence, $(f \star g) \star h = (f \star h) \star g$.

Theorem 2.2. $(T_l(X), \star, 0_X), (T_c(X), \star, 0_X), (T_r(X), \star, 0_X)$, and $(T(X), \star, 0_X)$ are BCH-algebras.

Let $X = (X, *, 0)$ be a BCH-algebra. We define the binary operation \diamond on X^3 as follows: $(\forall (a, b, c), (x, y, z) \in X^3)((a, b, c) \diamond (x, y, z) = (a * x, b * y, c * z))$. Then $X^3 = (X, \diamond, (0, 0, 0))$ is a BCH-algebra.

Theorem 2.3. Let $X = (X, *, 0)$ be a BCH-algebra and S_1, S_2, S_3 be subsets of X . Then

- (1) $S_1 \times S_2 \times S_3$ is a subalgebra of X^3 if and only if S_1, S_2 and S_3 are subsets of X ,
- (2) $S_1 \times S_2 \times S_3$ is an ideal of X^3 if and only if S_1, S_2 and S_3 are ideals of X .

Proof. (1) Suppose that $S_1 \times S_2 \times S_3$ is a subalgebra of X^3 . Firstly, we will show that S_1 is a subalgebra of X . Let $a, b \in S_1$. Let $x \in S_2$ and $u \in S_3$. Then $(a, x, u), (b, x, u) \in S_1 \times S_2 \times S_3$. Thus $(a * b, 0, 0) = (a * b, x * x, u * u) = (a, x, u) \diamond (b, x, u) \in S_1 \times S_2 \times S_3$, that is, $a * b \in S_1$. Hence, S_1 is a subalgebra of X . On the other hand, we can show that S_2 and S_3 are subalgebras of X .

Conversely, let $(x, y, z), (a, b, c) \in S_1 \times S_2 \times S_3$. Then $x * a \in S_1, y * b \in S_2$, and $z * c \in S_3$, so $(x, y, z) \diamond (a, b, c) = (x * a, y * b, z * c) \in S_1 \times S_2 \times S_3$. Hence, $S_1 \times S_2 \times S_3$ is a subalgebra of X^3 .

(2) Suppose that $S_1 \times S_2 \times S_3$ is an ideal of X^3 . Since $(0, 0, 0) \in S_1 \times S_2 \times S_3$, we have $0 \in S_i$ for all $i = 1, 2, 3$. Assume that $a * x \in S_1$ and $a \in S_1$. Let $b \in S_2$ and $c \in S_3$. Then $(a, b, c) \in S_1 \times S_2 \times S_3$ and $(x, b, c) \in X^3$. Thus $(a, b, c) \diamond (x, b, c) = (a * x, b * b, c * c) = (a * x, 0, 0) \in S_1 \times S_2 \times S_3$. Since $S_1 \times S_2 \times S_3$ is an ideal of X^3 , we have $(x, b, c) \in S_1 \times S_2 \times S_3$, that is, $x \in S_1$. Hence, S_1 is an ideal of X . Similarly, we can show that S_2 and S_3 are ideals of X .

Conversely, suppose that S_1, S_2 and S_3 are ideals of X . Since $0 \in S_i$ for all $i = 1, 2, 3$, we have $(0, 0, 0) \in S_1 \times S_2 \times S_3$. Assume that $(a, b, c) * (x, y, z) \in S_1 \times S_2 \times S_3$ and $(a, b, c) \in S_1 \times S_2 \times S_3$. We get $(a * x, b * y, c * z) \in S_1 \times S_2 \times S_3$. Since $a * x, a \in S_1$, we have $x \in S_1$. Moreover, we can obtain that $y \in S_2$ and $z \in S_3$. This implies that $(x, y, z) \in S_1 \times S_2 \times S_3$. Hence, $S_1 \times S_2 \times S_3$ is an ideal of X^3 . \square

Acknowledgment: This research project was supported by the Thailand Science Research and Innovation Fund and the University of Phayao (Grant No. FF66-UoE017).

Conflicts of Interest: The authors declare that there are no conflicts of interest regarding the publication of this paper.

References

- [1] R.K. Bandru, N. Rafi, On G -algebras, Sci. Magna, 8 (2012), 1-7.
- [2] W.A. Dudek, J. Thomys, On decompositions of BCH-algebras, Math. Japon. 35 (1990), 1131-1138. <https://cir.nii.ac.jp/crid/1572261549626076288>.
- [3] Q.P. Hu, X. Li, On BCH-algebras, Math. Seminar Notes, 11 (1983), 313-320.
- [4] K. Iséki, An algebra related with a propositional calculus, Proc. Japan Acad. Ser. A Math. Sci. 42 (1966), 26-29. <https://doi.org/10.3792/pja/1195522171>.
- [5] K. Iséki, On BCI-algebras, Math. Seminar Notes, 8 (1980), 125-130.

Taylor & Francis Journals

Uncover Breakthroughs in Pharmaceutical Science and Medical Research with our Key Titles

Taylor and Francis

Open

International Journal of Analysis and Applications

COUNTRY

Canada



Universities and research institutions in Canada



Media Ranking in Canada

SUBJECT AREA AND CATEGORY

Business, Management and Accounting
Business and International Management

Mathematics
Analysis
Applied Mathematics
Geometry and Topology

PUBLISHER

Semnan University, Center of Excellence in Nonlinear Analysis and Applications

H-INDEX

5

PUBLICATION TYPE

Journals









ISSN

22918639

COVERAGE

2021-2022

Join the conversation about this journal




Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimaç
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2023

Ebrima Bittaye 2 months ago

I want to publish my paper to International Journal of Analysis and Applications

reply



Melanie Ortiz 2 months ago

SCImago Team

Dear Ebrima,
Thank you for contacting us.
We suggest you visit the journal's homepage (See submission/author guidelines) or contact the journal's editorial staff , so they could inform you more deeply.
Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)



ฉันไม่ใช่โปรแกรมอัตโนมัติ

reCAPTCHA
มีอยู่ส่วนบุคคล - ชื่อกำหนด

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1, 1, 106)

[Cookie settings](#)

[Cookie policy](#)